

ON AMENABILITY OF GROUP ALGEBRAS, II: GRADED ALGEBRAS

LAURENT BARTHOLDI

ABSTRACT. We show that, in a finitely generated amenable group G with lower central series $(\gamma_n(G))$, the function $n \mapsto \text{rank}(\gamma_n(G)/\gamma_{n+1}(G))$ grows subexponentially.

This paper continues [22, 2]’s study of amenability of affine algebras (based on the notion of almost-invariant finite-dimensional subspace), and applies it to graded algebras associated with finitely generated groups.

We consider the graded deformation associated with the filtration of $\mathbb{k}G$ by powers of its augmentation ideal, and show that it has subexponential growth if G is amenable. This yields the statement in the first paragraph, and answers a question by Vershik [2424, 24], and another one by de la Harpe [1414, 14].

We also consider the graded deformation of a group ring $\mathbb{k}G$ associated with a metric on G , and note that this deformation is amenable whenever G is — but also if G has “dead ends” in its Cayley graph.

Warning! This paper contains a mistake in Lemma 4.3, and its main claims should not be considered as proven!

1. INTRODUCTION

Throughout this paper, let $\mathbb{k}$ denote a commutative ring, and for a $\mathbb{k}$ -module M let $\text{rank}(M)$ denote its minimal number of generators. Recall from [22, 2] the notion of amenable algebra, which first appeared (in a slightly different form) in [66, 61111, 11] and in [2323, 23] in the context of C^* -algebras:

Definition 1.1. Let R be a $\mathbb{k}$ -algebra and let M be a right R -module. It is *amenable* if, for every $\epsilon > 0$ and every finite-rank subspace S of R , there exists a finite-rank subspace F of M such that

$$\frac{\text{rank}(F + FS) - \text{rank}(F)}{\text{rank}(F)} < \epsilon;$$

any such F is called (S, ϵ) -invariant.

M is *exhaustively amenable* if furthermore in the definition above the space F may be required to contain any specified finite-rank subspace.

This is a counterpart to amenability of G -sets for a group G , where according to Følner’s definition [99, 9] the G -set X is *amenable* if, for every $\epsilon > 0$ and every finite subset $S \subset G$, there exists a finite subset $F \subset X$ such that $(\#(F \cup FS) - \#F)/\#F < \epsilon$; such F being also called (S, ϵ) -invariant. The main result of [22, 2] was:

Theorem 1.2. *Let $\mathbb{K}$ be a field, and let X be a G -set. Then X is amenable if and only if its linear envelope $\mathbb{K}X$ is amenable.*

1.1. Growth of modules. Let R be an *affine* algebra, i.e. a finitely generated associative algebra over a commutative ring $\mathbb{k}$, generated by the finite-rank subspace S . Let M be a finitely-generated R -module, generated (as an R -module) by a finite-rank subspace T . Let us write here and below

$$TS^{\leq n} = T + TS + TS^2 + \cdots + TS^n.$$

The module M is then filtered by the exhausting sequence of subspaces $T \leq TS^{\leq 1} \leq TS^{\leq 2} \leq \cdots \leq TS^{\leq n} \leq \cdots$, and the *growth* of M is the sequence $(r_n)_{n \geq 1}$ defined by $r_n = \text{rank}(TS^{\leq n})$. We say that M has *subexponential growth* if $\lim \sqrt[n]{r_n} = 1$. This property does not depend on the choices of S and T .

Proposition 1.3. *If M has subexponential growth, then M is exhaustively amenable.*

Proof. Given $U \leq M$ of finite rank and $\epsilon > 0$, let $d \in \mathbb{N}$ be such that $U \leq TS^{\leq d}$. Since (r_n) grows subexponentially, $\liminf r_{n+1}/r_n = 1$, so there exists $n > d$ such that $r_{n+1}/r_n < 1 + \epsilon$. Set $F = TS^{\leq n}$. Then $F + FS \leq TS^{\leq n+1}$, so $(\text{rank}(FS) - \text{rank}(F))/\text{rank}(F) < \epsilon$ and F contains U . $\square$

We consider in §2 the graded module $M_0 = \bigoplus_n TS^{\leq n}/TS^{\leq n-1}$ associated with the ascending filtration of M by its finite-rank subspaces TS^n . If we consider $R = \mathbb{k}G$ a group ring, S the $\mathbb{k}$ -span of a finite generating set of G , and $M = R$ with $T = \{1\}$, then $M_0 = R_0$ is an algebra with basis in bijection with G , and with product derived from multiplication in G except that $g \cdot h = 0$ if the length of gh is strictly less than the sum of the lengths of g and h . We prove:

Theorem 1.4. *Let $G = \langle S \rangle$ be a finitely generated group. If G is amenable, or has dead ends, then $(\mathbb{k}G)_0$ is amenable. If G is amenable, or has infinitely many dead ends, then $(\mathbb{k}G)_0$ is exhaustively amenable.*

The proof of Theorem 1.4 relies on fact that $(\mathbb{k}G)_0$ has a monomial basis, namely a basis B with $B \cdot B \subseteq B \cup \{0\}$, this prompts the

Question 1.5. *If M is an amenable finitely-generated R -module, is then M_0 also amenable?*

1.2. Augmented algebras. Assume now that R has an augmentation¹ $\varepsilon : R \rightarrow \mathbb{k}$, with kernel ϖ ; as a typical example $R = \mathbb{k}G$ and $\varpi = \langle g - 1 : g \in G \rangle$. Let M be an R -module. Then M admits a descending filtration $M \geq M\varpi \geq M\varpi^2 \geq \cdots \geq M\varpi^n \geq \cdots$; and an associated graded module

$$\overline{M} = \bigoplus_{n \geq 0} M\varpi^n / M\varpi^{n+1}.$$

In particular, R admits an associated graded algebra $\overline{R} = \bigoplus_{n \geq 0} \varpi^n / \varpi^{n+1}$. If M is generated (as an R -module) by a subspace T , then $\overline{M}$ is a graded $\overline{R}$ -module, generated by the image $\overline{T}$ of T in $M/\varpi M$.

If R is affine, say generated by a finite-rank subspace S , then $\overline{R}$ is also affine: the projection $\overline{S} = \{s - \varepsilon(s) + \varpi^2 : s \in S\}$ of S in ϖ/ϖ^2 actually equals ϖ/ϖ^2 irrespective of S , and generates $\overline{R}$. The *growth* of $\overline{M}$ is then its growth in the sense of §1.1, with respect to the generating subspace $\overline{S}$ of $\overline{R}$ and $\overline{T}$ of $\overline{M}$.

The main result of this paper is the following theorem, proven in §5:

¹Namely, a morphism of unital algebras to the scalars $\mathbb{k}$.

Theorem 1.6. *If G is an amenable, finitely generated group, then $\overline{\mathbb{k}G}$ has subexponential growth (and therefore is exhaustively amenable by Proposition 1.3).*

Vershik conjectured in [2424; 24, page 326] that if G is an amenable, finitely generated group and if ϖ denote the augmentation ideal in $\mathbb{Z}G$, then $\text{rank}(\varpi^n/\varpi^{n+1})$ grows subexponentially. This follows from Theorem 1.6 with $\mathbb{k} = \mathbb{Z}$.

Theorem 1.6 actually follows from the following statement: if R is an amenable augmented affine algebra with a basis consisting of invertible elements, then $\overline{R}$ has subexponential growth. This prompts the

Question 1.7. *Let R be an amenable augmented affine algebra. Does $\overline{R}$ necessarily have subexponential growth?*

Let more generally M be an amenable R -module, where R is any augmented affine algebra. Does $\overline{M}$ necessarily have subexponential growth?

1.3. Golod-Shafarevich groups. Let $\mathbb{K}$ be a field of characteristic $p > 0$, and assume that G is a residually- p group. This means that the series $(G_{n,p})_{n \geq 1}$ of dimension subgroups defined by $G_{1,p} = G$ and $G_{n+1,p} = [G_{n,p}, G](G_{[n/p],p})^p$ for $n \geq 1$, satisfies $\bigcap G_{n,p} = \{1\}$. The degree of $g \in G$, written $\deg_p(g)$, is the maximal $n \in \mathbb{N} \cup \{\infty\}$ such that $g \in G_{n,p}$.

It is well known [1717, 171919, 19] that $\mathcal{L} := \bigoplus_{n \geq 1} (G_{n,p}/G_{n+1,p}) \otimes_{\mathbb{F}_p} \mathbb{K}$ has the structure of a restricted Lie algebra over $\mathbb{K}$; and that $\overline{\mathbb{K}G}$ is the universal restricted enveloping algebra of $\mathcal{L}$. We will use this fact to relate Theorem 1.6 to purely group-theoretical statements, e.g. Corollary 1.11.

Golod constructed in [1010, 10] for every prime p a finitely generated infinite torsion p -group. His method is quite flexible, and was generalized as follows:

Definition 1.8. A group G is a *Golod-Shafarevich group* for the prime p if it admits a presentation $G = F/\langle \mathcal{R} \rangle^F$ in which F is a free group of rank d and $\mathcal{R} \subset F$ is a set of relators, such that for some $t \in (0, 1)$ we have

$$1 - dt + \sum_{r \in \mathcal{R}} t^{\deg_p r} < 0,$$

where $\deg_p$ denotes degree with respect to the filtration $(F_{n,p})_{n \geq 1}$ of F .

Proposition 1.9 ([1515, 15]; [1818, 18]; [1616; 16, §VIII.12]; [11, 1]). *All Golod-Shafarevich groups are infinite. If ϖ denote the augmentation ideal in $\mathbb{F}_p G$, then $\dim_{\mathbb{F}_p}(\varpi^n/\varpi^{n+1})$ grows exponentially (at rate at least $1/t$).*

On the other hand, there are torsion groups that are Golod-Shafarevich (this solved Burnside's problem [55, 5]).

The second part of the following result answers [1414; 14, Open Problem 5.2] and [1313; 13, Question 7]:

Corollary 1.10. *Golod-Shafarevich groups are not amenable. In particular, there exist non-amenable residually- p torsion groups.*

Ershov [77, 7] has constructed Golod-Shafarevich groups that have property (T) — and no infinite (T) group can be amenable. Any Golod-Shafarevich group admits a Golod-Shafarevich torsion quotient, which will still have property (T). This answers [1414; 14, Open Problem 5.2] by a different method.

1.4. Lower central series. Theorem 1.6 has the following purely group-theoretic consequence, also proven in §5:

Corollary 1.11. *Let G be an amenable, finitely-generated group, and let $(\gamma_n(G))_{n \geq 1}$ denote its lower central series. Then the function $n \mapsto \text{rank}(\gamma_n(G)/\gamma_{n+1}(G))$ grows subexponentially.*

Note that this function may grow arbitrarily close to an exponential function. Indeed Petrogradsky showed in [2121, 21] that if G be the free k -generated soluble group of solubility class $q \geq 3$, then

$$\text{rank}(\gamma_n(G)/\gamma_{n+1}(G)) \cong \exp \left(\left(\frac{(k-1)\zeta(k)}{\log \log \cdots n} \right)^{1/k} n \right),$$

with $q-3$ iterated logarithms in the expression above.

The converse of Theorem 1.6 does not hold (see Remark 5.4). However, the next statement trivially follows from Theorem 1.6, and raises the question after it:

Corollary 1.12. *Let G be an amenable group. Then for every finitely-generated subgroup H of G the function $n \mapsto \text{rank}(\gamma_n(H)/\gamma_{n+1}(H))$ grows subexponentially.*

Question 1.13. *Does there exist a non-amenable residually-nilpotent² group G such that for every finitely-generated subgroup H of G the function $n \mapsto \text{rank}(\gamma_n(H)/\gamma_{n+1}(H))$ grows subexponentially?*

1.5. Acknowledgments. The author is grateful to Mikhail Ershov, François Guéritaud, Pierre de la Harpe and Fabrice Krieger for generous feedback and/or entertaining and stimulating discussions.

2. HECKE AND CRYSTAL ALGEBRAS

We prove Theorem 1.4 in this section, phrasing it in a slightly more general manner. Let G be a group with fixed generating set S . Denote by $\ell(g)$ the length of $g \in G$ in the word metric:

$$\ell(g) = \min\{n : g = s_1 \dots s_n, s_i \in S\}.$$

Choose $\lambda \in \mathbb{k}$, and define the “Hecke algebra”³ $(\mathbb{k}G)_\lambda$ as follows: it is isomorphic to $\mathbb{k}G$ as a $\mathbb{k}$ -module; it has a basis $\{\delta_g\}_{g \in G}$; and multiplication is given by

$$\delta_g \delta_h = \lambda^{\ell(g)+\ell(h)-\ell(gh)} \delta_{gh}.$$

Note that, although the notation does not make it explicit, $(\mathbb{k}G)_\lambda$ depends on the choice of S .

If λ is invertible, then $(\mathbb{k}G)_\lambda$ is isomorphic to $\mathbb{k}G$, the isomorphism being given by $\delta_g \mapsto \lambda^{\ell(g)} g$. The universal cases $\mathbb{k} = \mathbb{Z}[\lambda]$ and $\mathbb{Q}[[\lambda]]$ should deserve particular consideration.

Quite on the contrary, $(\mathbb{k}G)_0$ is a graded algebra (with degree function ℓ), which we call the *crystal*⁴ of $\mathbb{k}G$. Note that $(\mathbb{k}G)_0$ is the associated graded algebra $\bigoplus \mathbb{k}S^{\leq n}/\mathbb{k}S^{\leq n-1}$ of the filtered algebra considered in §1.1. The first part of Theorem 1.4 generalizes as:

²Namely, such that $\bigcap \gamma_n(G) = 1$.

³The terminology comes from the classical Hecke algebra associated with the symmetric group.

⁴The terminology comes from statistical mechanics, where the parameter λ of the deformation of $\mathbb{k}G$ is interpreted as temperature.

Proposition 2.1. *Let G be an amenable group with fixed generating set. Then $(\mathbb{k}G)_\lambda$ is amenable for all $\lambda \in \mathbb{k}$.*

Proof. Let $\epsilon > 0$ be given, and let⁵ S' be a finite-rank subspace of $(\mathbb{k}G)_\lambda$. Let S denote the support of S' , i.e. the set of those $g \in G$ such that δ_g has a non-zero coefficient in some element of S' ; it is a finite subset of G . Since G is amenable, there exists a finite subset F of G with

$$(2.1) \quad (\#(F \cup FS) - \#F) / \#F < \epsilon.$$

Set $F' = \bigoplus_{g \in F} \mathbb{k}\delta_g$, a finite-rank subspace of $(\mathbb{k}G)_\lambda$. We have $\text{rank } F' = \#F$ and $F'S' \leq \mathbb{k}(FS)$, so $\text{rank}(F'S') \leq \#FS$, and $\text{rank}(F' + F'S') \leq \#(F \cup FS)$, whence

$$(2.2) \quad \frac{\text{rank}(F' + F'S') - \text{rank}(F')}{\text{rank}(F')} < \epsilon,$$

so $(\mathbb{k}G)_\lambda$ is amenable. $\square$

It is however possible for $(\mathbb{k}G)_0$ to be amenable, yet for G not to be amenable. The example in Proposition 2.3 appears in [44, 4] (with a small typographical mistake).

2.1. Dead ends. Say $g \in G$ is a *dead end* if $\ell(gs) \leq \ell(g)$ for all $s \in S$. Note that, although the notation does not make it apparent, this property strongly depends on the choice of S . Zoran Šuník has informed me that every group admits a generating set for which the group contains a dead end. The second part of Theorem 1.4 reads:

Proposition 2.2. *If G has a dead end (with respect to S), then $(\mathbb{k}G)_0$ is amenable. If G has infinitely many dead ends, then $(\mathbb{k}G)_0$ is exhaustively amenable.*

Proof. Let first $g \in G$ be a dead end. Set $F = \mathbb{k}\delta_g$. Then $F + Fs = F$ for any $s \in (\mathbb{k}G)_0$, so $(\mathbb{k}G)_0$ is amenable.

Let now $g_1, g_2, \dots$ be an infinite set of dead ends in G . Given $\epsilon > 0$ and $E \leq (\mathbb{k}G)_0$ of rank n , consider $F = E + \mathbb{k}\delta_{g_1} + \dots + \mathbb{k}\delta_{g_{\lceil n/\epsilon \rceil}}$. Then F contains E and $\text{rank}(F + Fs) \leq \text{rank } F + \text{rank } E$ and $\text{rank } F \geq \text{rank } E/\epsilon$. $\square$

Proposition 2.3 (Bogopolski). *For all $k \geq 3$, the “triangle” group*

$$G_k = T_{3,3,k} = \langle x, y \mid x^3, y^3, (xy)^k \rangle$$

contains infinitely many dead ends.

Proof. This group is hyperbolic; it acts by isometries on hyperbolic space and preserves the semiregular tiling of $\mathbb{H}^2$ by triangles (with edges labeled x and y) and $2k$ -gons (with edges labeled periodically x, y). The 1-skeleton of this tiling is the Cayley graph of G_k .

Assume first that k is even. Consider for all $n \in \mathbb{Z} \setminus \{0\}$ the element $d_n \in G_k$ defined by

$$d_{2n} = ((xy)^{k/2}(yx)^{k/2})^n, \quad d_{2n+1} = ((xy)^{k/2}(yx)^{k/2})^n(xy)^{k/2}.$$

Consider furthermore the automorphism ϕ of G_k defined by $x \mapsto y^{-1}$ and $y \mapsto x^{-1}$. It is easy to check that d_n is fixed by ϕ .

Geometrically, d_n is the hyperbolic isometry whose axis cuts through a doubly-infinite sequence of vertex-abutting $2k$ -gons in their middle, and translates by n of them; ϕ is the reflection through this axis.

⁵In this proof, we use S for a subset of a group, and S' for a subspace of an algebra.

Let $s_1 \dots s_m$ be a word of minimal length representing d_n , with $s_i \in S = \{x, y, x^{-1}, y^{-1}\}$; so $\ell(d_n) = m$. Then $\ell(d_n s) \leq m$ for all $s \in S$:

- if $s = s_m^{-1}$ this is clear;
- if $s = s_m$ then $d_n s = s_1 \dots s_{m-1} s_m^{-1}$ whence $\ell(d_n s) \leq m$;
- otherwise, $d_n^\phi = d_n = s_1^\phi \dots s_m^\phi$ also of minimal length, and $s_m^\phi \in \{s, s^{-1}\}$, so the previous cases apply.

If k is odd, consider for all $n \neq 0$ the element $d_n = ((xy)^{(k-1)/2}x)^n$, with a similar geometric interpretation as above. The same arguments apply. $\square$

It is in fact not hard to see that these are the only dead ends in G_k . Since for $k \geq 4$ the G_k are non-elementary hyperbolic groups, they are certainly not amenable.

There unfortunately does not seem to be any natural condition to impose on $(\mathbb{k}G)_0$ to ensure that G be amenable.

3. TILEABLE AMENABLE GROUPS

We prove in this section a result by Weiss [2525, 25], based on earlier work by Ornstein and Weiss [2020; 20, §I.2]. We follow the sketch of a proof by Gromov [1111; 11, pages 336–337], adapting it so as to prepare the ground for a generalization to modules in §4. We will not use the results in this §, but present them as a warm-up for §4.

Let X denote a G -set. We defined amenability with respect to the *outer envelope* AK of a subset $A \subset X$ with respect to $K \subset G$. It will be useful in this section to consider, again for $A \subset X$ and $K \subset G$, the *inverse envelope*

$$AK^* := \{x \in X : xK \cap A \neq \emptyset\}.$$

The easy properties

$$A(K \cup L)^* = AK^* \cup AL^*, \quad (A \cup B)K^* = AK^* \cup BK^*$$

follow immediately from $AK^* = A\{k^{-1} : k \in K\}$.

Theorem 3.1 (Weiss). *Let G be an amenable group, let $K \subseteq G$ be a finite subset, let $\epsilon > 0$ be given, and let $N_0 \geq N_1 \geq \dots$ be a nested sequence of finite-index normal subgroups of G such that $\bigcap_{n \in \mathbb{N}} N_n = \{1\}$.*

Then for all $n \gg 0$ there exists a (K, ϵ) -invariant subset $T_n \subseteq G$ that is a transversal for N_n in G .

Let $\zeta \geq 1$ and $\delta \in (0, 1)$ be constants to be fixed later. In this section and the next we shall use, for $\mu \geq \delta$, the transformation $\Theta_\mu : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+^2$ given by

$$\Theta_\mu(\nu, \alpha) = \left(\nu + \mu(1 - \alpha), \alpha + \frac{\mu(1 - \alpha)}{1 - \delta} \zeta \right).$$

The proof of Theorem 3.1 relies on the following

Lemma 3.2. *Let Ω be a finite G -set; let B be a subset of Ω and let K, L be finite subsets of G ; assume that for all $x \in \Omega$ the orbit map*

$$(3.1) \quad K \rightarrow \Omega, \quad k \mapsto xk \quad \text{is injective.}$$

Let $\alpha, \nu \in [0, 1)$ and $\zeta \geq 1$ be such that

$$\begin{aligned} \#(KL^*) &\leq \zeta \#K, & \#(BK^*) &\leq \alpha \# \Omega, \\ \#B &= \nu \# \Omega, & \#(BL^*) &\leq \alpha \# \Omega; \end{aligned}$$

Let furthermore $\delta \in (0, 1)$ be given.

Then there exist $s \geq 1$, elements $x_1, \dots, x_s \in \Omega$, and $\mu \geq \delta$ such that, setting $B_0 = B$, $B_i = B_{i-1} \cup x_i K$ for $1 \leq i \leq s$ and $\Theta_\mu(\nu, \alpha) = (\nu', \alpha')$, we have

$$(3.2) \quad \#(x_i K \cap B_{i-1}) \leq \delta \#K \text{ for all } i \in \{1, \dots, s\};$$

$$(3.3) \quad \#B_s = \nu' \# \Omega;$$

$$(3.4) \quad \#(B_s L^*) \leq \alpha' \# \Omega.$$

Informally, the lemma says that, given a set K with small L -boundary, and a set $B \subset \Omega$ with small K - and L -boundary, one can construct a quantifiably larger set B_s with small L -boundary.

Proof. Let $x_1, \dots, x_s \in \Omega$ be a maximal-length sequence of elements such that (3.2) holds. Then, by maximality of $(x_1, \dots, x_s)$, for all $x \in \Omega$ we have

$$(3.5) \quad \#(B_s \cap xK) = \#\{(b, k) : b \in B_s, k \in K, xk = b\} > \delta \#K.$$

We deduce

$$\begin{aligned} \#B_s \#K &= \#\{(b, k, x) : b \in B_s, k \in K, x \in \Omega, xk = b\} \\ &= \sum_{x \in \Omega} \#\{(b, k) : b \in B_s, k \in K, xk = b\} = \sum_{x \in \Omega} \#(B_s \cap xK) \\ &= \sum_{x \in BK^*} \#\{(b, k) : b \in B_s, k \in K, xk = b\} + \sum_{x \in \Omega \setminus BK^*} \#(B_s \cap xK) \\ &= \sum_{b \in B_s} \sum_{k \in K} \#\{BK^* \cap \{bk^{-1}\}\} + \sum_{x \in \Omega \setminus BK^*} \#(B_s \cap xK) \\ &\geq \sum_{b \in B} \sum_{k \in K} \#\{BK^* \cap \{bk^{-1}\}\} + \sum_{x \in \Omega \setminus BK^*} \delta \#K \text{ by (3.5)} \\ &\geq \#B \#K + (1 - \alpha) \# \Omega \delta \#K. \end{aligned}$$

If we divide by $\#K \# \Omega$ and set $\mu = (\#B_s / \# \Omega - \nu) / (1 - \alpha)$, then we get $\mu \geq \delta$, and therefore $s \geq 1$. Note that (3.3) holds by the choice of μ . Next,

$$\#B_s \geq \#B + \sum_{i=1}^s \#(x_i K \setminus B_{i-1}) \geq \#B + s(1 - \delta) \#K \text{ by (3.1, 3.2)}$$

so $s \#K \leq (\nu' - \nu) \# \Omega / (1 - \delta)$; and

$$\#(B_s L^*) \leq \#(B L^*) + \sum_{i=1}^s \#(x_i K L^*) \leq \alpha \# \Omega + s \zeta \#K,$$

from which (3.4) follows. $\square$

Remark 3.3. We shall apply Lemma 3.2 using the following strategy: we construct a finite but very long sequence $K_1, \dots, K_t$ of Følner sets, each with a very small boundary with respect to its predecessors and to K . We then find a finite quotient Ω of G in which these Følner sets embed. Then, starting from K_t down to K_1 , we apply t times Lemma 3.2 to cover most of Ω by images of translates of the K_i , taking each time as many copies as possible subject to them having extremely small overlaps (see condition (3.2)). There remains a small part of Ω that is not covered. We then lift these images back to G , and lift the small remainder arbitrarily. We




FIGURE 1. The proof of Theorem 3.1, illustrated in a picture

have obtained a transversal, consisting mostly of pieces carved out of Følner sets by other Følner sets. See Figure 1 for an illustration.

Note that both coördinates of Θ_μ increase monotonically in μ . This property helps us formalize the intuitive notion that “it cannot hurt us, in constructing the covering of Ω , if the biggest Følner sets leave unexpectedly small holes between them, even though small holes are harder to cover efficiently with smaller Følner sets”. $\triangle$

Proof of Theorem 3.1. We are given K and ϵ . Without loss of generality, we suppose $1 \in K$. Fix in order the following data:

- (1) Choose $\delta > 0$, the “allowed overlap” as in (3.2), such that $\delta \#K < \frac{\epsilon}{2}$ and $(1 + \frac{\epsilon}{2})(1 - \delta) > 1$;
- (2) Choose $\zeta > 1$ such that $(1 - \delta)/\zeta > 1 - \epsilon/(2\#K)$, the “relative Følner constant”.

For all $t \in \mathbb{N}$ set $(\overline{\nu}_t, \overline{\alpha}_t) = \Theta_\delta^t(0, 0)$. Then it is clear that $\overline{\nu}_t/\overline{\alpha}_t = (1 - \delta)/\zeta$ for all $t > 0$, and that $\limsup \overline{\alpha}_t \geq 1$, so $\limsup \overline{\nu}_t \geq (1 - \delta)/\zeta$.

- (3) Let $t \in \mathbb{N}$, the “tower height”, be such that $\overline{\nu}_t > 1 - \epsilon/(2\#K)$.

Then for any $\mu_1, \dots, \mu_t \geq \delta$ we also have

$$\Theta_{\mu_t}(\dots \Theta_{\mu_1}(0, 0) \dots) \in (1 - \epsilon/(2\#K), 1] \times [0, 1].$$

- (4) Using amenability of G , construct finite subsets $K_0 = K, K_1, \dots, K_t$ of G , the “Rokhlin tower”, such that for all $j < i$ in $\{1, \dots, t\}$ one has $\#(K_i K_j^*) < \zeta \#K_i$, and such that

$$(3.6) \quad \#(K_i K) \leq (1 + \frac{\epsilon}{2})(1 - \delta) \#K_i \text{ for all } i \in \{1, \dots, t\}.$$

- (5) Choose $n \in \mathbb{N}$, the “quotient index”, large enough so that $K_i K_i^* \cap N_n = \{1\}$ for all $i \in \{1, \dots, t\}$.

Set $\Omega = G/N_n$, let $\pi : G \rightarrow \Omega$ denote the natural quotient map, and remark, by the choice of n , that $K_i \rightarrow \Omega$, $k \mapsto xk$ is injective for all $i \in \{1, \dots, t\}$ and all $x \in \Omega$.

Now start with $A_{t,0} = \emptyset$ and $(\nu_t, \alpha_t) = (0, 0) = (\overline{\nu}_0, \overline{\alpha}_0)$, and apply t times Lemma 3.2: at step $i = t, t-1, \dots, 1$, apply it with $B := A_{i,0}$, $K := \pi(K_i)$, $L := \pi(K_{i-1})$, and $(\nu, \alpha) := (\nu_i, \alpha_i)$. Lemma 3.2 produces for us

- constants $s \geq 1$ and $\mu \geq \delta$, which we rechristen $s(i)$ and μ_i ;
- a sequence $x_1, \dots, x_{s(i)}$ in Ω , which we rechristen $\overline{x}_{i,1}, \dots, \overline{x}_{i,s(i)}$;
- a sequence $B_0 \subseteq \dots \subseteq B_{s(i)}$ of subsets of Ω , which we rechristen $A_{i,0}, \dots, A_{i,s(i)} =: A_{i-1,0}$.

Set $(\nu_{i-1}, \alpha_{i-1}) = \Theta_{\mu_i}(\nu_i, \alpha_i)$, and note that $\nu_{i-1} \geq \overline{\nu}_{t-(i-1)}$ by induction on i and because $\mu_i \geq \delta$ and Θ_μ is monotonically increasing in μ . Finally define $K_{i,j} \subset G$ by $\overline{x}_{i,j} K_{i,j} = \overline{x}_{i,j} K_i \setminus A_{i,j-1}$, and note by (3.2) that

$$(3.7) \quad \#(\overline{x}_{i,j} K_{i,j}) \geq (1 - \delta) \#K_i.$$

If $\alpha_{i-1} \geq 1$, then for all $j < i-1$ set $A_{j,0} = A_{i-1,0}$, $s(j) = 0$ and $\nu_j = \nu_{i-1}$, and stop; otherwise, decrease i and continue.

After all these steps, we have obtained a decomposition

$$(3.8) \quad \Omega = \overline{Q} \sqcup \bigsqcup_{i=1}^t \bigsqcup_{j=1}^{s(i)} \overline{x}_{i,j} K_{i,j},$$

with $\overline{Q} = \Omega \setminus A_{0,0}$. If the iteration was stopped because $\alpha_{i-1} \geq 1$, then $\nu_0 = \alpha_{i-1}(1 - \delta)/\zeta > 1 - \epsilon/(2\#K)$; otherwise, $\nu_0 \geq \overline{\nu}_t > 1 - \epsilon/(2\#K)$. In all cases, we have

$$(3.9) \quad \#\overline{Q} \leq (1 - \nu_0) \# \Omega < \frac{\epsilon}{2\#K} \# \Omega.$$

Lift $\overline{Q}$ and $\overline{x_{i,j}}$ to $Q \subseteq G$ and $x_{i,j} \in G$ respectively. We have obtained a finite subset

$$(3.10) \quad T_n = Q \sqcup \bigsqcup_{i=1}^t \bigsqcup_{j=1}^{s(i)} x_{i,j} K_{i,j}$$

of G . Furthermore, the natural restriction $\pi|_{T_n}: T_n \rightarrow \Omega$ is a bijection, i.e. T_n is a transversal to N_n . We compute

$$\begin{aligned} \#(T_n \cup T_n K) &= \#(T_n K) \text{ since } 1 \in K \\ &\leq \#(QK) + \sum_{i,j} \#(x_{i,j} K_i K) \text{ by (3.10)} \\ &\leq \#Q \#K + \sum_{i,j} (1 + \frac{\epsilon}{2})(1 - \delta) \#(x_{i,j} K_i) \text{ by (3.6)} \\ &\leq \#Q \#K + \sum_{i,j} (1 + \frac{\epsilon}{2}) \#(x_{i,j} K_{i,j}) \text{ by (3.7)} \\ &\leq \frac{\epsilon \# \Omega}{2 \# K} \#K + (1 + \frac{\epsilon}{2}) \# \Omega \text{ by (3.9 + 3.8)} \\ &\leq (1 + \epsilon) \# T_n. \end{aligned} \quad \square$$

4. TILEABLE AMENABLE ALGEBRAS

We prove in this section an analogue of Theorem 3.1 for algebras. We follow as much as possible the notation of the previous section; this mainly amounts to replacing ‘ $\#$ ’ by ‘ $\dim$ ’, ‘ $\sqcup$ ’ by ‘ $\oplus$ ’ and so on. The equation numbers match between these two sections; Section 3 also contains a few informal remarks that may help the reader along the proof(s).

Throughout this section we consider an associative algebra R over a field $\mathbb{K}$, we use $\oplus$ and $\otimes$ to denote direct sum and tensor products as $\mathbb{K}$ -vector spaces, and we denote by ‘ $\dim$ ’ the dimension as a $\mathbb{K}$ -vector space. An *i-subspace* of R is a subspace of R admitting a basis consisting of invertible elements. An *i-algebra* is an algebra R which is an i-subspace of itself, and R is *i-amenable* if for every finite-dimensional subspace K and every $\epsilon > 0$ there exists a (K, ϵ) -invariant i-subspace of R , in the sense of an i-subspace F such that

$$\frac{\text{rank}(F + FK) - \text{rank}(F)}{\text{rank}(F)} < \epsilon.$$

Lemma 4.1. *The group ring of an amenable group is an i-amenable i-algebra.*

Proof. The basis G of $\mathbb{K}G$ consists of invertible elements. Following the proof of Proposition 2.1, there exist (K, ϵ) -invariant subspaces of the form $\mathbb{K}F$ with $F \subseteq G$, which have a basis F consisting of invertible elements. $\square$

Given a finite-dimensional subspace A and a finite-dimensional i-subspace K of R , with fixed basis $\{k_1, \dots, k_d\} \subseteq R^\times$, define

$$AK^* := A \mathbb{K}\{k_1^{-1}, \dots, k_d^{-1}\}.$$

Note that, although the notation does not make it apparent, AK^* may depend on the choice of basis of K .

Theorem 4.2. *Let R be an amenable i -algebra, let $K \leq R$ be a finite-dimensional i -subspace, let $\epsilon > 0$ be given, and let $I_0, I_1, \dots$ be a sequence of finite-codimension ideals in R such that $\bigcap_{n \in \mathbb{N}} I_n = \{0\}$.*

Then for all $n \gg 0$ there exists a (K, ϵ) -invariant subspace $T_n \leq R$ that is a vector-space complement for I_n in R .

Let $\zeta \geq 1$ and $\delta \in (0, 1)$ be constants to be fixed later. In this section we shall use, for $\mu \geq \delta$, the transformation $\Theta_\mu : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+^2$ given by

$$\Theta_\mu(\nu, \alpha) = \left(\nu + \mu(1 - \alpha), \alpha + \frac{\mu(1 - \alpha)}{1 - \delta} \zeta \right).$$

The proof of Theorem 4.2 relies on the following

Lemma 4.3. *Let Ω be a finite-dimensional R -module over an i -algebra R , with spanning subset $\mathcal{C}$; let B be a subspace of Ω and let K, L be finite-dimensional i -subspaces of R ; assume that for all $x \in \mathcal{C}$ the orbit map*

$$(4.1) \quad K \rightarrow \Omega, \quad k \mapsto xk \quad \text{is injective.}$$

Let $\alpha, \nu \in [0, 1)$ and $\zeta \geq 1$ be such that

$$\begin{aligned} \dim(KL^*) &\leq \zeta \dim K, & \dim(BK^*) &\leq \alpha \dim \Omega, \\ \dim B &= \nu \dim \Omega, & \dim(BL^*) &\leq \alpha \dim \Omega; \end{aligned}$$

Let furthermore $\delta \in (0, 1)$ be given.

Then there exist $s \geq 1$, elements $x_1, \dots, x_s \in \mathcal{C}$, and $\mu \geq \delta$ such that, setting $B_0 = B$ and $B_i = B_{i-1} + x_i K$ for $1 \leq i \leq s$ and $\Theta_\mu(\nu, \alpha) = (\nu', \alpha')$, we have

$$(4.2) \quad \dim(x_i K \cap B_{i-1}) \leq \delta \dim K \text{ for all } i \in \{1, \dots, s\};$$

$$(4.3) \quad \dim B_s = \nu' \dim \Omega;$$

$$(4.4) \quad \dim(B_s L^*) \leq \alpha' \dim \Omega.$$

Proof. Let $x_1, \dots, x_s \in \mathcal{C}$ be a maximal-length sequence of elements such that (4.2) holds. Let $\{v_1, \dots, v_d\}$ be a maximal subset of $\mathcal{C}$ whose image is independent in Ω/BK^* ; then $V := \mathbb{k}\{v_1, \dots, v_d\}$ is a vector space complement to BK^* in Ω . By maximality of $(x_1, \dots, x_s)$, for all $x \in \mathcal{C}$ we have

$$(4.5) \quad \dim(B_s \cap xK) = \dim \mathbb{k}\{x \otimes k \in \Omega \otimes K : xk \in B_s\} > \delta \dim K.$$

Let K have i -basis $\{k_i\} \subseteq R^\times$; then for any $A \leq \Omega$ the vector space $\{\sum \omega_i \otimes k_i \in \Omega \otimes K : \omega_i k_i \in A \text{ for all } i\}$ is isomorphic to $A \otimes K$ via the map

$$(4.†) \quad \sum \omega_i \otimes k_i \mapsto \sum \omega_i k_i \otimes k_i$$

with inverse $\sum \xi_i \otimes k_i \mapsto \sum \xi_i k_i^{-1} \otimes k_i$. We compute

$$\begin{aligned} \dim B_s \dim K &= \dim(B_s \otimes K) \\ &= \dim \left\{ \sum \omega_i \otimes k_i \in \Omega \otimes K : \omega_i \in B_s \forall i \right\} \\ &= \dim \left\{ \sum \omega_i \otimes k_i \in \Omega \otimes K : \omega_i k_i \in B_s \forall i \right\} \text{ by (4.†)} \\ &\geq \dim \left\{ \sum \omega_i \otimes k_i \in BK^* \otimes K : \omega_i k_i \in B_s \forall i \right\} \end{aligned}$$

$$\begin{aligned}
& + \dim \left\{ \sum \omega_i \otimes k_i \in V \otimes K : \omega_i k_i \in B_s \forall i \right\} \\
& \geq \dim \left\{ \sum \omega_i k_i^{-1} \otimes k_i \in BK^* \otimes K : \omega_i \in B_s \forall i \right\} + \dim(VK \cap B_s) \\
& \geq \dim(B \otimes K) + \sum_{j=1}^d \dim(v_j K \cap B_s) \text{ by (4.5)} \\
& \geq \dim B \dim K + d\delta \dim K;
\end{aligned}$$

and $d \geq (1 - \alpha) \dim \Omega$. If we divide by $\dim K \dim \Omega$ and set, $\mu = (\dim B_s / \dim \Omega - \nu) / (1 - \alpha)$, then we get $\mu \geq \delta$, and therefore $s \geq 1$. Note that (4.3) holds by the choice of μ . Next,

$$\begin{aligned}
\dim B_s & \geq \dim B + \sum_{i=1}^s \dim(x_i K / (x_i K \cap B_{i-1})) \\
& \geq \dim B + s(1 - \delta) \dim K \text{ by (4.1, 4.2),}
\end{aligned}$$

so $s \dim K \leq (\nu' - \nu) \dim \Omega / (1 - \delta)$; and

$$\dim(B_s L^*) \leq \dim(BL^*) + \sum_{i=1}^s \dim(x_i K L^*) \leq \alpha \dim \Omega + s\zeta \dim K,$$

from which (4.4) follows. $\square$

For a sketch of the proof of Theorem 4.2, which applies as well to algebra setting as to the group setting, see Remark 3.3 on page 7.

Proof of Theorem 4.2. Without loss of generality (possibly adjoining a unit to R first), we suppose $1 \in K$. Fix in order the following data:

- (1) Choose $\delta > 0$, the “allowed overlap” as in (4.2), such that $\delta \dim K < \frac{\epsilon}{2}$ and $(1 + \frac{\epsilon}{2})(1 - \delta) > 1$;
- (2) Choose $\zeta > 1$ such that $(1 - \delta)/\zeta > 1 - \epsilon/(2 \dim K)$, the “relative Følner constant”.

For all $t \in \mathbb{N}$ set $(\overline{\nu}_t, \overline{\alpha}_t) = \Theta_\delta^t(0, 0)$. Then it is clear that $\overline{\nu}_t / \overline{\alpha}_t = (1 - \delta)/\zeta$ for all $t > 0$, and that $\limsup \overline{\alpha}_t \geq 1$, so $\limsup \overline{\nu}_t \geq (1 - \delta)/\zeta$.

- (3) Let $t \in \mathbb{N}$, the “tower height”, be such that $\overline{\nu}_t > 1 - \epsilon/(2 \dim K)$.

Then for any $\mu_1, \dots, \mu_t \geq \delta$ we also have

$$\Theta_{\mu_t}(\dots \Theta_{\mu_1}(0, 0) \dots) \in (1 - \epsilon/(2 \dim K), 1] \times [0, 1].$$

- (4) Using i-amenability of R , construct finite-dimensional i-subspaces $K_0 = K, K_1, \dots, K_t$ of R , the “Rokhlin tower”, such that for all $j < i$ in $\{1, \dots, t\}$ one has $\dim(K_i K_j^*) < \zeta \dim K_i$, and such that

$$(4.6) \quad \dim(K_i K) \leq \left(1 + \frac{\epsilon}{2}\right)(1 - \delta) \dim K_i \text{ for all } i \in \{1, \dots, t\}.$$

- (5) Choose $n \in \mathbb{N}$, the “quotient index”, large enough so that $K_i K_i^* \cap I_n = (0)$ for all $i \in \{1, \dots, t\}$.

Set $\Omega = G/I_n$, let $\pi : R \rightarrow \Omega$ denote the natural quotient map, and let $\mathcal{C}$ denote the image in Ω of the spanning set $R^\times$ of R . Remark, by the choice of n , that $K_i \rightarrow \Omega, k \mapsto xk$ is injective for all $i \in \{1, \dots, t\}$ and all $x \in \mathcal{C}$.

Now start with $A_{t,0} = \{0\} \leq \Omega$ and $(\nu_t, \alpha_t) = (0, 0) = (\overline{\nu_0}, \overline{\alpha_0})$, and apply t times Lemma 4.3: at step $i = t, t-1, \dots, 1$, apply it with $B := A_{i,0}$, $K := \pi(K_i)$, $L := \pi(K_{i-1})$, and $(\nu, \alpha) := (\nu_i, \alpha_i)$. Lemma 4.3 produces for us

- constants $s \geq 1$ and $\mu \geq \delta$, which we rechristen $s(i)$ and μ_i ;
- a sequence $x_1, \dots, x_{s(i)}$ in $\mathcal{C}$, which we rechristen $\overline{x_{i,1}}, \dots, \overline{x_{i,s(i)}}$;
- a sequence $B_0 \subseteq \dots \subseteq B_{s(i)}$ of subspaces of Ω , which we rechristen $A_{i,0}, \dots, A_{i,s(i)} =: A_{i-1,0}$.

Set $(\nu_{i-1}, \alpha_{i-1}) = \Theta_{\mu_i}(\nu_i, \alpha_i)$, and note that $\nu_{i-1} \geq \overline{\nu_{t-(i-1)}}$ by induction on i and because $\mu_i \geq \delta$ and Θ_μ is monotonically increasing in μ . Finally let $K_{i,j} \leq K_i$ be any subspace such that $\overline{x_{i,j}}K_{i,j} \oplus (\overline{x_{i,j}}K_i \cap A_{i,j-1}) = \overline{x_{i,j}}K_i$, and note by (4.2) that

$$(4.7) \quad \dim(\overline{x_{i,j}}K_{i,j}) \geq (1 - \delta) \dim K_i,$$

because the $\overline{x_{i,j}}$ are invertible.

If $\alpha_{i-1} \geq 1$, then for all $j < i-1$ set $A_{j,0} = A_{i-1,0}$, $s(j) = 0$ and $\nu_j = \nu_{i-1}$, and stop; otherwise, decrease i and continue.

After all these steps, we have obtained a decomposition

$$(4.8) \quad \Omega = \overline{Q} \oplus \bigoplus_{i=1}^t \bigoplus_{j=1}^{s(i)} \overline{x_{i,j}}K_{i,j},$$

where $\overline{Q}$ is any vector space complement to $A_{0,0}$ in Ω . If the iteration was stopped because $\alpha_{i-1} \geq 1$, then $\nu_0 = \alpha_{i-1}(1 - \delta)/\zeta > 1 - \epsilon/(2 \dim K)$; otherwise, $\nu_0 \geq \overline{\nu_t} > 1 - \epsilon/(2 \dim K)$. In all cases, we have

$$(4.9) \quad \dim \overline{Q} \leq (1 - \nu_0) \dim \Omega < \frac{\epsilon}{2 \dim K} \dim \Omega.$$

Lift $\overline{Q}$ and $\overline{x_{i,j}}$ to $Q \leq R$ and $x_{i,j} \in R^\times$ respectively. We have obtained a finite-dimensional subspace

$$(4.10) \quad T_n = Q \oplus \bigoplus_{i=1}^t \bigoplus_{j=1}^{s(i)} x_{i,j}K_{i,j}$$

of R . Furthermore, the natural restriction $\pi|_{T_n}: T_n \rightarrow \Omega$ is a bijection, i.e. T_n is a vector space complement to I_n . We compute

$$\begin{aligned} \dim(T_n + T_n K) &= \dim(TK) \text{ because } 1 \in K \\ &\leq \dim(QK) + \sum_{i,j} \dim(x_{i,j}K_i K) \text{ by (4.10)} \\ &\leq \dim Q \dim K + \sum_{i,j} (1 + \frac{\epsilon}{2})(1 - \delta) \dim(x_{i,j}K_i) \text{ by (4.6)} \\ &\leq \dim Q \dim K + \sum_{i,j} (1 + \frac{\epsilon}{2}) \dim(x_{i,j}K_i) \text{ by (4.7)} \\ &\leq \frac{\epsilon \dim \Omega}{2 \dim K} \dim K + (1 + \frac{\epsilon}{2}) \dim \Omega \text{ by (4.9 + 4.8)} \\ &\leq (1 + \epsilon) \dim T_n. \end{aligned}$$

□

5. PROOF OF THEOREM 1.6 AND COROLLARY 1.11

We start by a “Reidemeister-Schreier” result for algebras and ideals:

Lemma 5.1. *Let R be a unital $\mathbb{k}$ -algebra generated by a subspace S ; let $I \triangleleft R$ be a right ideal in R ; and let $F \leq R$ be a complement of I , so we have $R = I \oplus_{\mathbb{k}} F$. Let $x \mapsto \bar{x}$ be the projection $R \rightarrow F$; assume that $1 \in F$ and $\bar{1} = 1$. Then*

$$I = \{fs - \overline{fs} : f \in F, s \in S\}R.$$

Proof. Write $J = \langle fs - \overline{fs} \rangle$; then obviously $J \subseteq I$. Conversely, consider first $s_1 \dots s_n \in R$, and write $\equiv_J$ for congruence modulo J . Then

$$\begin{aligned} s_1 \dots s_n &= (\bar{1}s_1 - \overline{s_1})s_2 \dots s_n + \overline{s_1}s_2 \dots s_n \\ &\equiv_J (\overline{s_1}s_2 - \overline{s_1s_2})s_3 \dots s_n + \overline{s_1s_2}s_3 \dots s_n \\ &\equiv_J \dots \equiv_J \overline{s_1 \dots s_n}. \end{aligned}$$

Consider now any $x = \sum_i s_{i,1} \dots s_{i,n_i} \in R$. Then $x \equiv_J \sum_i \overline{s_{i,1} \dots s_{i,n_i}} \in F$, so $R = J + F$ and therefore $I = J$. $\square$

Remark 5.2. Consider the right ideal $I = (H - 1)\mathbb{k}G \triangleleft \mathbb{k}G$ for some subgroup H of $G = \langle S \rangle$. Let T be a right transversal of H in G ; then $\mathbb{k}T$ is a complement of I in $\mathbb{k}G$, so I is generated by $\{fs - \overline{fs} : f \in T, s \in S\}$ by Lemma 5.1, and therefore also by $\{fs\overline{fs}^{-1} - 1\}$; so H is generated by $\{fs\overline{fs}^{-1} : s \in S, f \in T\}$, which is the Reidemeister-Schreier generating set of H . $\triangle$

Corollary 5.3. *Let $R = \langle S \rangle$ be an augmented algebra, and consider $I \triangleleft R$ with $R = I \oplus_{\mathbb{k}} F$. Then $I/I\varpi$ is spanned by the image of $(F + FS) \cap I$ in $I/I\varpi$.*

We are now ready to prove Theorem 1.6. We start by a special case:

Proof of Theorem 1.6 for $\mathbb{k} = \mathbb{K}$ a field of positive characteristic. Let a finite subset S of G and $\eta > 1$ be given; we will show that $\lim \sqrt[n]{\dim(\mathbb{K}G/\varpi^n)} \leq \eta$.

Since $\mathbb{K}$ has positive characteristic and G is finitely generated, the dimension subgroups $G_n = (1 + \varpi^n) \cap G$ have finite index in G for all n . Since any quotient of an amenable group is amenable, we may replace G by $G/\bigcap_{n \geq 1} G_n$ and assume from now on that $\bigcap G_n = 1$. The ideals ϖ^n have finite codimension in $\mathbb{K}G$, and have trivial intersection.

We apply Theorem 4.2 to $I_n = \varpi^n$: let $n_0 \in \mathbb{N}$ be such that for all $n \geq n_0$ there is a subspace F_n of $\mathbb{K}G$ with $\mathbb{K}G = I_n \oplus_{\mathbb{K}} F_n$ and $\dim(F_n + F_n S) < \eta \dim F_n$. Then

$$\begin{aligned} \dim(\mathbb{K}G/\varpi^{n+1}) &= \dim(\mathbb{K}G/\varpi^n) + \dim(\varpi^n/\varpi^{n+1}) \\ &= \dim F_n + \dim(I_n/I_n\varpi) \\ &\leq \dim F_n + \dim((F_n + F_n S) \cap I_n) \text{ by Corollary 5.3} \\ &\leq \dim((F_n + F_n S) \cap F_n) + \dim((F_n + F_n S) \cap I_n) \\ &\leq \dim((F_n + F_n S) \cap (F_n + I_n)) = \dim(F_n + F_n S) \\ &\leq \eta \dim F_n = \eta \dim(\mathbb{K}G/\varpi^n). \end{aligned}$$

Set $C = \dim(\mathbb{K}G/\varpi^{n_0})/\eta^{n_0}$. We therefore have $\dim(\mathbb{K}G/\varpi^n) < C\eta^n$ for all $n \geq n_0$, so $\lim \sqrt[n]{\dim(\mathbb{K}G/\varpi^n)} \leq \eta$ for all $\eta > 1$. $\square$

Proof of Theorem 1.6 for $\mathbb{k} = \mathbb{Z}$. For $n \geq 0$ set $r_n = \text{rank}(\varpi^n/\varpi^{n+1})$, where ϖ denotes the augmentation ideal of $\mathbb{Z}G$; for $n \geq 0$ and p prime set $s_{n,p} = \text{rank}(\varpi_p^n/\varpi_p^{n+1})$, where ϖ_p denotes the augmentation ideal of $\mathbb{F}_pG$; let $r_{n,p} = \text{rank}(\varpi^n/\varpi^{n+1} \otimes_{\mathbb{Z}} \mathbb{Z}[\frac{1}{p}])$ denote the rank of the part of ϖ^n/ϖ^{n+1} that is coprime to p , and let $r_{n,0} = \text{rank}(\varpi^n/\varpi^{n+1} \otimes_{\mathbb{Z}} \mathbb{Q})$ denote the free rank of ϖ^n/ϖ^{n+1} .

Since G is finitely generated (say by d elements), ϖ^n/ϖ^{n+1} is a finite-rank abelian group (of rank at most d^n), so $\cong \mathbb{Z}^{r_{n,0}} \oplus \text{torsion}$. We thus have $r_{n,p} \geq r_{n,0}$, and for fixed n we have $r_{n,p} = r_{n,0}$ for almost all p .

By Theorem 1.6 for $\mathbb{k} = \mathbb{F}_p$, the sequence $s_{n,p}$ grows subexponentially for any fixed p . Every $\mathbb{F}_p$ factor in $\varpi_p^n/\varpi_p^{n+1}$ lifts to a $\mathbb{Z}$ -factor in $\mathbb{Z}G$, which then gives either a $\mathbb{Z}$ -factor in ϖ^n/ϖ^{n+1} , or gives a torsion factor in ϖ^m/ϖ^{m+1} for all $m \geq n$. Therefore $r_{m,p} \leq \sum_{n \leq m} s_{n,p}$, so $r_{n,p}$ grows subexponentially for any fixed p .

The multiplication maps $\varpi^m/\varpi^{m+1} \otimes \varpi^n/\varpi^{n+1} \rightarrow \varpi^{m+n}/\varpi^{m+n+1}$ are onto, so the sequence r_n is submultiplicative ($r_m r_n \geq r_{m+n}$); the same holds for the sequences $r_{n,p}$ for fixed p , and for the sequence $r_{n,0}$.

Let $\eta > 1$ be given. Then for some $n \in \mathbb{N}$ we have $r_{n,0} < \eta^n$; and for some p_0 we have $r_{n,p} = r_{n,0}$ if $p > p_0$. By submultiplicativity, $r_{kn,p} \leq r_{n,p}^k < \eta^{kn}$ for all k .

For all $p \leq p_0$ there exists $k_p \in \mathbb{N}$ such that $r_{k_p n, p} < \eta^{k_p n}$, because $r_{n,p}$ grows subexponentially. Set $m = n \cdot \text{lcm}(k_1, \dots, k_{p_0})$. Then $r_{m,p} \leq \eta^m$ for all p .

We have $r_n = \max_p r_{n,p}$ for all $n \in \mathbb{N}$, so $r_{km} \leq \eta^{km}$ for all $k \in \mathbb{N}$. Since r_n is submultiplicative, we have

$$\limsup \sqrt[n]{r_n} = \lim (r_{kn})^{1/kn} \leq \eta$$

by Fekete's Lemma [2222; 22, volume 1, part I, problem 98; originally [88; 8, page 233]]. Since $\eta > 1$ was arbitrary, the sequence r_n grows subexponentially. $\square$

Proof of Theorem 1.6 for general $\mathbb{k}$. Let ϖ denote the augmentation ideal of $\mathbb{Z}G$, and let $\overline{\varpi}$ denote the augmentation ideal of $\mathbb{k}G$. Since the natural map $\varpi^n/\varpi^{n+1} \otimes \mathbb{k} \rightarrow \overline{\varpi}^n/\overline{\varpi}^{n+1}$ is onto for all $n \in \mathbb{N}$, we have $\text{rank}_{\mathbb{k}}(\mathbb{k}G/\overline{\varpi}^n) \leq \text{rank}_{\mathbb{Z}}(\mathbb{Z}G/\varpi^n)$, so the claim follows from Theorem 1.6 for $\mathbb{k} = \mathbb{Z}$. $\square$

Remark 5.4. The converse of Theorem 1.6 does not hold: the group $SL(d, \mathbb{Z})$ for $d \geq 3$ is certainly not amenable (it contains free subgroups), and neither is its congruence subgroup $K = \ker(SL(d, \mathbb{Z}) \rightarrow SL(d, \mathbb{Z}/p\mathbb{Z}))$. This subgroup is residually- p if $p \geq 3$, since⁶ the subgroups $K_n = \ker(SL(d, \mathbb{Z}) \rightarrow SL(d, \mathbb{Z}/p^n\mathbb{Z}))$ have trivial intersection and index $p^{(n-1)(d^2-1)}$ in K . Then, because K has the congruence property [33, 3], the congruence subgroup K_n coincides with the dimension subgroup as defined in §1.2, so $\dim(\varpi^n/\varpi^{n+1})$ grows subexponentially (approximately at rate $e^{(d^2-1)\pi\sqrt{2n/3}}$). I wish to thank M. Ershov for pointing out this example to me. $\triangle$

Corollary 1.11 could follow along the same lines as the proof for $\mathbb{k} = \mathbb{Z}$ of Theorem 1.6, by reducing from $\gamma_n(G)/\gamma_{n+1}(G)$ to quotients of p -dimension subgroups $G_{n,p}/G_{n+1,p}$ at all primes p , and using the fact that $\bigoplus_{n \geq 1} G_{n,p}/G_{n+1,p}$ is the primitive part of the Hopf algebra $\overline{\mathbb{F}_pG}$ and therefore has subexponential growth. We will however opt for a shortcut:

⁶As usual in these situations, one should treat sometimes 2, sometimes 4 as the even prime to extend this result to characteristic 2.

Proof of Corollary 1.11. The classical *dimension subgroups* of G are the subgroups $\delta_n(G) = G \cap (1 + \varpi^n)$, where ϖ denotes the augmentation ideal in $\mathbb{Z}G$. By a result of Gupta [1212, 12], the quotient $\delta_n(G)/\gamma_n(G)$ is a finite 2-group. Now $\text{rank}(\delta_n(G)/\delta_{n+1}(G))$ grows subexponentially by Theorem 1.6 for $\mathbb{k} = \mathbb{Z}$, since $\delta_n(G)/\delta_{n+1}(G)$ is a submodule of ϖ^n/ϖ^{n+1} ; and $\dim_{\mathbb{F}_2} G_{n,2}/G_{n+1,2}$ grows subexponentially since $\overline{\mathbb{F}_2 G}$ has subexponential growth by Theorem for $\mathbb{k} = \mathbb{F}_2$. We conclude that

$$\begin{aligned} \text{rank}(\gamma_n(G)/\gamma_{n+1}(G)) &\leq \text{rank}(\gamma_n(G)/\gamma_{n+1}(G) \otimes \mathbb{Z}[\tfrac{1}{2}]) + \text{rank}(\gamma_n(G)/\gamma_{n+1}(G) \otimes \mathbb{F}_2) \\ &\leq \text{rank}(\delta_n(G)/\delta_{n+1}(G)) + \text{rank}(G_{n,2}/G_{n+1,2}) \end{aligned}$$

grows subexponentially. $\square$

REFERENCES

- [1] [cite.bartholdi-g:lie1Laurent Bartholdi and Rostislav I. Grigorchuk, *Lie methods in growth of groups and groups of finite width*, Computational and geometric aspects of modern algebra (Edinburgh, 1998), London Math. Soc. Lecture Note Ser., vol. 275, Cambridge Univ. Press, Cambridge, 2000, pp. 1–27, available at [arXiv.org/abs/math/0002010](https://arxiv.org/abs/math/0002010). MR **1776763** (2001h:20046)
- [2] [cite.bartholdi:aa12Laurent Bartholdi, *On amenability of group algebras, I* (2007), to appear in *Isr. J. Math.*, available at [arXiv.org/abs/math/0608302](https://arxiv.org/abs/math/0608302).
- [3] [cite.bass-l-s:congruence3Hyman Bass, Michel Lazard, and Jean-Pierre Serre, *Sous-groupes d'indice fini dans $\mathbf{SL}(n, \mathbf{Z})$* , Bull. Amer. Math. Soc. **70** (1964), 385–392 (French). MR 0161913 (28 #5117)
- [4] [cite.bogopolski:bilipschitz4Oleg V. Bogopol'skiĭ, *Infinite commensurable hyperbolic groups are bi-Lipschitz equivalent*, Algebra i Logika **36** (1997), no. 3, 259–272, 357 (Russian, with Russian summary); English transl., Algebra and Logic **36** (1997), no. 3, 155–163. MR **1485595** (98h:57002)
- [5] [cite.burnside:question5William Burnside, *On an unsettled question in the theory of discontinuous groups*, Quart. J. Pure Appl. Math. **33** (1902), 230–238.
- [6] [cite.elek:amena6Gábor Elek, *The amenability of affine algebras*, J. Algebra **264** (2003), no. 2, 469–478. MR **1981416** (2004d:16043)
- [7] [cite.erшов:goshat7Mikhail V. Eršov, *Golod-Shafarevich groups with property (T) and Kac-Moody groups* (2006). Preprint.
- [8] [cite.fekete:verteilung8Mihály Fekete, *Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten*, Math. Z. **17** (1923), no. 1, 228–249 (German). MR 1544613
- [9] [cite.følner:banach9Erling Følner, *Note on groups with and without full Banach mean value*, Math. Scand. **5** (1957), 5–11.
- [10] [cite.golod:nil10Evgeniĭ S. Golod, *On nil-algebras and finitely approximable p -groups*, Izv. Akad. Nauk SSSR Ser. Mat. **28** (1964), 273–276.
- [11] [cite.gromov:topinv11Mikhael L. Gromov, *Topological invariants of dynamical systems and spaces of holomorphic maps. I*, Math. Phys. Anal. Geom. **2** (1999), no. 4, 323–415. MR **1742309** (2001j:37037)
- [12] [cite.gupta:odddimension12Narain D. Gupta, *The dimension subgroup conjecture holds for odd order groups*, J. Group Theory **5** (2002), no. 4, 481–491. MR **1931371** (2003m:20019)
- [13] [cite.harpe:mfp13Pierre de la Harpe, *Mesures finiment additives et paradoxes*, Autour du centenaire Lebesgue, Panor. Synthèses, vol. 18, Soc. Math. France, Paris, 2004, pp. 39–61 (French, with English and French summaries). MR **2143416** (2006e:43001)
- [14] [cite.harpe:uniform14Pierre de la Harpe, *Uniform growth in groups of exponential growth*, Geom. Dedicata **95** (2002), 1–17.
- [15] [cite.herstein:rings15Israel N. Herstein, *Noncommutative rings*, Carus Mathematical Monographs, vol. 15, Mathematical Association of America, Washington, DC, 1994. Reprint of the 1968 original; With an afterword by Lance W. Small.

- [16] [cite.huppert-b:fg216]Bertram Huppert and Norman Blackburn, *Finite groups. II*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 242, Springer-Verlag, Berlin, 1982. AMD, 44. MR **650245** (**84i**:20001a)
- [17] [cite.jennings:gpring17]Stephen A. Jennings, *The structure of the group ring of a p -group over a modular field*, Trans. Amer. Math. Soc. **50** (1941), 175–185.
- [18] [cite.koch:galois18]Helmut Koch, *Galoissche Theorie der p -Erweiterungen*, Springer-Verlag, Berlin, 1970. Mit einem Geleitwort von I. R. Šafarevič.
- [19] [cite.lazard:uea19]Michel Lazard, *Sur les algèbres enveloppantes universelles de certaines algèbres de Lie*, C. R. Acad. Sci. Paris Sér. I Math. **234** (1952), 788–791.
- [20] [cite.ornstein-weiss:entropyiso20]Donald S. Ornstein and Benjamin Weiss, *Entropy and isomorphism theorems for actions of amenable groups*, J. Analyse Math. **48** (1987), 1–141. MR **910005** (**88j**:28014)
- [21] [cite.petrogradsky:polynilpotent21]Victor M. Petrogradskiĭ, *Growth of finitely generated polynilpotent Lie algebras and groups, generalized partitions, and functions analytic in the unit circle*, Internat. J. Algebra Comput. **9** (1999), no. 2, 179–212.
- [22] [cite.polya-s:analysis22]George Polyá and Gabor Szegő, *Aufgaben und Lehrsätze aus der Analysis. Band I: Reihen. Integralrechnung. Funktionentheorie*, Dritte berichtigte Auflage. Die Grundlehren der Mathematischen Wissenschaften, Band 19, Springer-Verlag, Berlin, 1964 (German). MR 0170985 (30 #1219a)
- [23] [cite.vaillant:folner23]Ghislain Vaillant, *Folner conditions, nuclearity, and subexponential growth in C^* -algebras*, J. Funct. Anal. **141** (1996), no. 2, 435–448. MR **1418514** (**97m**:46098)
- [24] [cite.vershik:amenability24]Anatoly M. Vershik, *Amenability and approximation of infinite groups*, Selecta Math. Soviet. **2** (1982), no. 4, 311–330. Selected translations. MR **721030** (**86g**:43006)
- [25] [cite.weiss:monotileable25]Benjamin Weiss, *Monotileable amenable groups*, Topology, ergodic theory, real algebraic geometry, Amer. Math. Soc. Transl. Ser. 2, vol. 202, Amer. Math. Soc., Providence, RI, 2001, pp. 257–262. MR **1819193** (**2001m**:22014)

ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE (EPFL), INSTITUT DE MATHÉMATIQUES B (IMB), 1015 LAUSANNE, SWITZERLAND
E-mail address: laurent.bartholdi@gmail.com